

## Atrovent® Lösung

### To whom it may concern:

The following description shall help patients to inform international authorities about their medication. It is not supposed to replace the original patient leaflet. Patients are recommended to keep it together with the medication and the original patient leaflet.

### Prescribed drug

<b>Trade name</b>	Atrovent® LS
<b>Dosage form:</b>	Solution for a nebulizer
<b>International name of active substance</b>	Ipratropium bromide
<b>Concentration of active substance</b>	1 ml solution, corresponding to 10 puffs from the dispenser, contains 0.25 mg ipratropium bromide  Excipients: 100 mcg benzalkonium chloride, sodium edetate, sodium chloride, hydrochloric acid 3.6%, purified water
<b>Indications</b>	For prophylaxis and treatment of dyspnoea associated with - chronic obstructive pulmonary disease (COPD) - mild to moderate bronchial asthma in adults and children as a supplement to beta-2-mimetics during acute asthma attack
<b>Dosage</b>	Dosage must be adapted to the individual requirements.  <b>For acute treatment:</b>  <u>Adults and adolescents over 12 years of age:</u> 20 puffs (0.5 mg ipratropium bromide); repeated doses can be administered until breathlessness (dyspnoea) has improved, time intervals between the single inhalations to be determined by a doctor.  <u>Children 6 - 12 years:</u> 10 puffs (0.25 mg ipratropium bromide); repeated doses can be administered until breathlessness (dyspnoea) has improved, time intervals between the single inhalations to be determined by a doctor.  <u>Children younger than 6 years:</u> 4 – 10 strokes (0.1 - 0.25 mg ipratropium bromide) under regular medical supervision; repeated doses can be administered until breathlessness (dyspnoea) has improved, time

	<p>intervals between the single inhalations to be determined by a doctor.</p> <p><b>For long-term treatment:</b></p> <p><u>Adults and adolescents over 12 years of age:</u> 20 puffs (0.5 mg ipratropium bromide) 3 to 4 times daily</p> <p><u>Children 6 - 12 years:</u> The single dose for inhalation is 0.25 mg ipratropium bromide, corresponding to 10 puffs; repeated doses can be administered until breathlessness (dyspnoea) has improved, time intervals between the single inhalations to be determined by a doctor.</p> <p><u>Children younger than 6 years:</u> The single dose for inhalation is 0.1 - 0.25 mg ipratropium bromide, corresponding to 4 – 10 puffs under regular medical supervision; repeated doses can be administered until breathlessness (dyspnoea) has improved, time intervals between the single inhalations to be determined by a doctor.</p>
<b>Available Sizes</b>	<p>N1    20 ml N3    5 x 20 ml</p>

Lieber Patient,

bitte beachten Sie, dass diese kurze Beschreibung ausschließlich dazu dient, bei Auslandsreisen die Behörden Ihres Einreiselandes über Art und Inhalt Ihres Arzneimittels zu informieren. Sie ersetzt nicht den Original-Beipackzettel. Wir empfehlen Ihnen, diese Beschreibung zusammen mit Ihrem Arzneimittel und dem Beipackzettel aufzubewahren und auf Verlangen vorzuzeigen.

Mit freundlichen Grüßen  
Boehringer Ingelheim Pharma GmbH & Co. KG