

Berodual® LS

To whom it may concern:

The following description shall help patients to inform international authorities about their medication. It is not supposed to replace the original patient leaflet. Patients are recommended to keep it together with the medication and the original patient leaflet.

Prescribed drug

Trade name	Berodual® LS 250 µg/500 µg solution for nebulizer
Dosage form:	Solution for nebulizer
International name of active substance	ipratropium bromide and fenoterol hydrobromide
Concentration of active substance	1 ml (=10 actuations with the solution dispenser) contains: ipratropium bromide 250 µg and fenoterol hydrobromide 500 µg
Indications	For prevention and treatment of shortness of breath in chronic obstructive airway disorders: allergic and non-allergic (endogenous) bronchial asthma, exercise-induced asthma and chronic obstructive bronchitis with and without emphysema. Note: If a long-term therapy is required, a concomitant anti-inflammatory therapy should be carried out always.
Dosage	The dosage depends on the nature and severity of the disease. Unless otherwise prescribed, the following dosages are recommended for adults and children: (the treatment should usually be started with the lowest recommended dose): <u>Adults and children over 12 years:</u> For acute treatment of sudden bronchial, depending on the severity of the acute episode, 10 – 25 actuations (1.0 – 2.5 ml) Berodual LS should be inhaled, after dilution with physiological saline to a volume of 3 – 4 ml. : In exceptional severe cases, up to 40 puffs (4 ml) Berodual LS may be inhaled, after dilution with physiological saline to 3 – 4 ml. For targeted prevention in exercise-induced asthma or predictable allergic contact, 1 – 2 puffs (0.1 – 0.2 ml) Berodual LS should be inhaled, diluted with 2 – 3 ml physiological saline,

	<p>, if possible 10 – 15 minutes before.</p> <p><u>Children 6 – 12 years:</u></p> <p>For acute treatment of acute asthma episodes or predictable allergic contact, depending on the severity of the acute episode and on age, 5 – 20 puffs (0.5 – 2.0 ml) Berodual LS should be inhaled, after dilution with physiological saline to a volume of 3 – 4 ml.</p> <p>For targeted prevention in exercise-induced asthma or predictable allergic contact, 1 – 2 puffs (0.1 – 0.2 ml) Berodual LS should be inhaled, diluted with 2 – 3 ml physiological saline, if possible 10 – 15 minutes before.</p> <p><u>Children under 6 years:</u></p> <p>Because there is limited information in this age group the following dose is recommended to be given under medical supervision only: 1 puff (0.1 ml) per kg body weight up to maximal 5 puffs (0.5 ml), after dilution with physiological saline to a volume of 3 – 4 ml.</p>
<p>Available Sizes in Germany</p>	<p>N1 20 ml N3 5 x 20 ml</p>

Lieber Patient,

bitte beachten Sie, dass diese kurze Beschreibung ausschließlich dazu dient, bei Auslandsreisen die Behörden Ihres Einreiselandes über Art und Inhalt Ihres Arzneimittels zu informieren. Sie ersetzt nicht den Original-Beipackzettel. Wir empfehlen Ihnen, diese Beschreibung zusammen mit Ihrem Arzneimittel und dem Beipackzettel aufzubewahren und auf Verlangen vorzuzeigen.

Mit freundlichen Grüßen
Boehringer Ingelheim Pharma GmbH & Co. KG